DEFENCE R&D DÉFENSE

Analysis of GA, GB, GD and GF in Aqueous Samples by Packed Capillary LC-ESI-MS

Paul A. D'Agostino, James R. Hancock and Lionel R. Provost
Defence Research Establishment Suffield
Box 4000, Stn Main
Medicine Hat, Alberta T1A 8K6

"The use of this information is permitted subject to recognition of proprietary and patent rights."

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD

Technical Memorandum
DRES TM 1999-047

February 1999

Défense nationale National Defence

Canada

102-02982/

UNCLASSIFIED

DRES Technical Memorandum 1999-047

Analysis of GA, GB, GD and GF in Aqueous Samples by Packed Capillary LC-ESI-MS

by

Paul A. D'Agostino, James R. Hancock and Lionel R. Provost

February 1999

UNCLASSIFIED

ABSTRACT

Packed capillary column liquid chromatography (LC) - electrospray mass spectrometry (ESI-MS) was used for the first time to detect and identify four common organophosphorus chemical warfare agents in aqueous samples. Aqueous samples containing the organophosphorus chemical warfare agents in the 0.01 to 0.1 mg/mL range were analysed directly by packed capillary LC-ESI-MS with the chemical warfare agents and several minor related impurities being well resolved under acetonitrile/water gradient elution conditions. The ESI-MS data for isopropyl methylphosphonofluoridate (GB or sarin), ethyl dimethylphosphoramidocyanidate (GA or tabun), cyclohexyl methylphosphonofluoridate (GF) and pinacolyl methylphosphonofluoridate (GD or soman) were acquired with a sampling cone voltage setting that promoted collisionally activated dissociation, and resulted in the acquisition of informative mass spectra containing both molecular and product ion information. The developed method appears to be an attractive alternative to GC-MS for the analysis of aqueous samples containing organophosphorus chemical warfare agents and their hydrolysis products, since they may be analysed directly without the need for additional sample handling.

Executive Summary

Title: P.A. D'Agostino, J.R. Hancock and L.R. Provost, "Analysis of GA, GB, GD and GF in Aqueous Samples by Packed Capillary LC-ESI-MS", Suffield Technical Memorandum No. 1999-047, 1999, UNCLASSIFIED.

Introduction: The Canadian Forces (CF) may be called on to perform peacekeeping or battlefield operations in regions of the world where there is a significant threat of chemical/biological (CB) warfare agent use. To operate effectively in these theatres the CF must be able to identify the CB agent used. Mass spectrometry (MS), is a powerful analytical technique for the identification of both known and unknown compounds and DRE Suffield is currently investigating this instrumental technique in fulfilment of CF detection and identification requirements.

Results: Packed capillary column liquid chromatography (LC) - electrospray mass spectrometry (ESI-MS) was used for the first time to detect and identify four common organophosphorus chemical warfare agents in aqueous samples. Aqueous samples containing the organophosphorus chemical warfare agents in the 0.01 to 0.1 mg/mL range were analysed directly by packed capillary LC-ESI-MS with the chemical warfare agents and several minor related impurities being well resolved under acetonitrile/water gradient elution conditions. The ESI-MS data for isopropyl methylphosphonofluoridate (GB or sarin), ethyl dimethylphosphoramidocyanidate (GA or tabun), cyclohexyl methylphosphonofluoridate (GF) and pinacolyl methylphosphono-fluoridate (GD or soman) were acquired with a sampling cone voltage setting that promoted collisionally activated dissociation, and resulted in the acquisition of informative mass spectra containing both molecular and product ion information. The developed method appears to be an attractive alternative to GC-MS for the analysis of aqueous samples containing organophosphorus chemical warfare agents and their hydrolysis products, since they may be analysed directly without the need for additional sample handling.

Significance of Results: The CF may be deployed in regions of the world where there is a significant threat of chemical/biological warfare agent use. Identification of the CB agent is of importance since the results of such analyses would contribute to the development of strategic and political positions regarding future Canadian military operations and would facilitate the dissemination of technical advice to in-theatre field commanders and medical personnel.

Future Goals: The reported ESI-MS data could prove valuable for the identification of organophosphorus chemical warfare agents in samples collected by the Canadian Forces, during base cleanup operations or in support of Chemical Weapons Convention challenge inspections.

TABLE OF CONTENTS

ABSTRACT..... i

EXECUTIVE SUMMARY.....ii

INTRODUCTION..... 1

EXPERIMENTAL..... 3

RESULTS AND DISCUSSION..... 5

CONCLUSIONS..... 8

REFERENCES..... 9

INTRODUCTION

The Chemical Weapons Convention entered into force almost two years ago, effectively banning the production, stockpiling and use of chemical weapons by all signatory nations. A strong compliance monitoring regime involving site inspections was built into the convention to ensure a verifiable treaty. Routine inspections have or will take place at declared sites, including small scale production, storage and destruction sites, and challenge inspection will take place at sites suspected of non-compliance. An analytical capability will be required to verify the convention, since inspectors will have the option to procure and analyse suspect samples to help establish compliance. Ongoing development of new, specific methods (1) for the detection and identification of chemical warfare agents, their degradation products and related compounds would benefit the inspectorate, as an improved analytical capability could act as an additional deterrent to non-compliance.

Gas chromatography (GC) has been used extensively for the separation and identification of the chemical warfare agents (1,2), with GC-MS being used most frequently for the characterization of these compounds (3-7). Organophosphorus chemical warfare agents, scheduled under the Chemical Weapons Convention, have been studied extensively by electron impact and chemical ionization mass spectrometry as the use of these complementary ionization techniques facilitates the acquisition of molecular and fragmentation ion information that may be used for unambiguous identification (8-10). GC separation, while generally suitable for the direct analysis of organophosphous chemical warfare agent in organic extracts, is usually not preferred for the direct analysis of aqueous samples. Aqueous samples containing organophosphorus chemical warfare agents and/or their nonvolatile hydrolysis products normally require additional sample handling steps and derivatization (11-13). Recently, water samples containing chemical warfare agents have been analysed by GC-MS following solid-phase microextraction (14) and by microcolumn liquid chromatography (LC) with flame photometric detection (15). Increasingly,

researchers have developed LC-MS separation methods to deal with the analysis of aqueous samples containing these nonvolatile hydrolysis products (16-21). Benefits over GC analysis include reduced or no sample handling and no requirement for derivatization to increase the volatility of the hydrolysis products.

Use of thermospray mass spectrometry (16-19) and more recently the atmospheric pressure ionization (e.g., electrospray (ESI), ionspray and atmospheric pressure CI) techniques (20-26) has enabled the direct mass spectrometric analysis of the hydrolysis products of organophosphorus chemical warfare agents. Both techniques may be interfaced to LC for component separation, with thermospray having been largely superseded by atmospheric pressure ionization (API) for most applications. Most recent API-MS and LC-API-MS papers have focussed on the analysis of the hydrolysis products of chemical warfare agents, with the exception of a recent presentation dealing with the direct aqueous LC-ESI-MS analysis of a degraded ethyl dimethylphosphoramidocyanidate sample (26). ESI-MS, the most sensitive technique for these applications (21), has not been previously used for the characterization of the organophosphorus chemical warfare agents, isopropyl methylphosphonofluoridate (GB or sarin), ethyl dimethylphosphoramidocyanidate (GA or tabun), cyclohexyl methylphosphonofluoridate (GF) or pinacolyl methylphosphonofluoridate (GD or soman). This paper focuses on the ESI-MS characterization of these four organophosphorus chemical warfare agents and the development of a LC-ESI-MS method for the direct analysis of these compounds in aqueous samples.

EXPERIMENTAL

Samples

GB, GA, GF and GD were synthesized and purified locally by the Organic Chemistry Laboratory at Defence Research Establishment Suffield. Stock solutions containing each of the four compounds were prepared in water (pH 5 to 6) at concentrations of 0.1mg/mL and 0.01 mg/mL.

Instrumental

All electrospray mass spectra were acquired using a Micromass Autospec-Q tandem mass spectrometer (Manchester, UK) equipped with the Mark II electrospray interface. The electrospray needle was operated at 7.6 kV and ions were accelerated into the mass spectrometer at 4 kV. Sampling cone voltages of 25 or 50 volts were utilized. Nitrogen (Very Dry, Liquid Carbonic Inc., Scarborough, Ont., Canada) bath gas was introduced into the interface (80 °C) at a flow rate of 300 L/hr. Nitrogen nebulizer gas was introduced at a flow rate of 14 L/hr. The electrospray interface was pumped with both a rotary and a turbomolecular pump, which enabled maintenance of a $4x10^{-4}$ and $7x10^{-6}$ Pa within the source and analyser regions of the instrument, respectively. LC-ESI-MS data were acquired in the continuum mode by scanning the magnetic sector from 600 to 60 u (7 sec/decade) with a resolution of 1000 (10% valley definition). Two to three scans were typically averaged to enhance the signal-to-noise ratio.

All LC separations were performed with an Applied Biosystems Model 140B dual syringe pump (Foster City, CA) equipped with a Zorbax 150 mm × 0.32 mm i.d. C_{18} SB (5 μm) packed fused-silica capillary column and a Rheodyne 8125 (Cotati, CA) injector with a 5 μL sample loop. The following solvent compositions were prepared for sample introduction: Solvent

A (0.1% trifluoroacetic acid in water) and Solvent B (0.1% trifluoroacidic acid in acetonitrile(ACN)/water, 95:5). Chromatographic separations were performed using a 1% to 75%B gradient program over 30 minutes. In order to minimize dead volume effects and ensure reproducible mixing, the mobile phase was delivered at 200 μL/min and split prior to the injector such that the flow through the column was 5 μL/min.

RESULTS AND DISCUSSION

GC-MS has used extensively for the detection and identification of organophosphorus chemical warfare agents in organic extracts (1,2), but this separation method does not generally permit direct analysis of aqueous samples containing organophosphorus chemical warfare agents and their hydrolysis products. The development of a complementary LC-MS method for these compounds would be beneficial as it would allow simultaneous identification of both organophosphorus chemical warfare agents and their hydrolysis products in a single direct analysis. LC-ESI-MS methods have been recently demonstrated for a series of alkyl methylphosphonic acid standards (21) and for the analysis of degradation products related to the hydrolysis of munitions grade mustard (27). Packed capillary column LC columns with an internal diameter of 0.32 mm were selected for this study and the prior mustard hydrolysis study since the 5 μL/min flow rates typically used during chromatographic separation with these packed capillaries approaches the lower flow rate limit for spraying in the ESI interface used. Optimal sensitivity resulted at this flow rate limit due to the concentration dependance of MS detection.

Chromatographic separation of the four organophosphorus chemical warfare agents was achieved using a 1% to 75%B gradient program over 30 minutes (Figure 1). The molecular masses of four organophosphorus chemical warfare agents, as well as some minor GA impurities, were established from the acquired ESI-MS data. These GA impurities, with retention times in the 12 to 14 minute range, were the same as several that have been reported previously during GC-MS study (3).

Figure 2 illustrates the ESI-MS data that were obtained for GB (molecular mass 140), GA (molecular mass 162), GF (molecular mass 180) and GD (molecular mass 182) with a sampling cone voltage of 25 volts. All the spectra exhibited $(M+H)^+$ and $(M+H+ACN)^+$ ions and

protonated dimers that could be used to confirm the molecular mass of each compound. Some additional minor adduct ions due to $(M+NH_4)^+$ and $(M+H_3O)^+$ were also generally observed (e.g., ions at m/z 158 and m/z 159 for GB (Figure 2a)).

Product ions due to alkene loss from the alkoxy substituents and the acetonitrile adduct associated with these product ions were observed for all four compounds. The ESI-MS data for GB (Figure 2a) contained a significant product ion at m/z 99 due to loss of C_3H_6 from the $(M+H)^+$ ion as well as its acetonitrile adduct at m/z 140. A product ion due to loss of C_2H_4 was observed for GA (Figure 2b) at m/z 135 along with its acetonitrile adduct at m/z 176. The ESI-MS data acquired for GF exhibited product ions at m/z 279 and m/z 99 (and its acetonitrile adduct at m/z 140) due to loss of the cyclic alkene, C_6H_{10}. GD was characterized by product ions due to loss of C_6H_{12} (and the associated acetonitrile adducts) from the protonated monomer and dimer at m/z 99 and m/z 281, respectively, and a product ion at m/z 85 due to $(C_6H_{13})^+$ and its associated acetonitrile adduct at m/z 126.

The formation of significant acetonitrile adducts was of initial concern. While this behaviour has been observed during some pharmaceutical analyses (28) it was not significant during a prior LC-ESI-MS study of mustard hydrolysis products (27). Adduct formation, similar to what has been observed for these compounds during ammonia CI-MS (9), was likely due to the lower proton affinity of the compounds coupled with the opportunity for multiple collisions within the ESI interface. Increasing the sampling cone voltage increased the relative abundance of the product ions due to alkene loss, but had little overall effect on the presence of acetonitrile adducts. The effect of the organic component in the mobile phase was then investigated by replacing the acetonitrile with methanol during gradient programming LC-ESI-MS analysis of a degraded sample of GA (26). The most notable change in the acquired mass spectra for GA and related compounds in the sample was the presence of methanol as opposed to acetonitrile adducts. Finally the presence of these solvent adducts was confirmed on a completely different

instrument, a Hewlett Packard 1100 MSD (quadrupole), equipped with an ESI interface of their own design.

A detailed detection limit study was not undertaken as the study focussed more on the separation and characterization of the organophosphorus chemical warfare agents. A full scanning (600 to 60 u) detection limit of 5 ng, based on the acquisition of an interpretable mass spectrum, was estimated during analysis of the 0.01 mg/mL standard mixture and a relatively pure GA standard with the same concentration (26). This limit was similar to that obtained for thiodiglycol, the hydrolysis product of mustard (27). Selected-ion-monitoring, which typically results in a 10 to 100 fold increase in sensitivity, was not evaluated.

CONCLUSIONS

This study represents the first application of packed capillary column LC-ESI-MS for the characterization of organophosphorus chemical warfare agents and demonstrates application of this method for the direct analysis of these compounds in aqueous samples. The ESI-MS data were collected with sampling cone voltages in the 25 to 50 volts range. In general the most informative mass spectra were acquired with the lower sampling cone voltage, a setting that promoted collisionally activated dissociation, and resulted in the acquistion of mass spectra containing both molecular and product ion information.

An LC-ESI-MS method has been demonstrated for the direct analysis of nanogram quantities of organophosphorus chemical warfare agents in aqueous samples, extending the range of analytical options available to the researcher confronted with the identification of chemical warfare agents or their hydrolysis products. The developed method appears to be an attractive alternative to GC-MS for the analysis of aqueous samples since they may be analysed directly reducing the need for additional sample handling or derivatization steps. Use of this method resulted in the ESI-MS characterization of GB, GA, GF and GD. The ESI-MS data generated during LC-ESI-MS analysis would be valuable during chemical weapons destruction monitoring of countries in compliance with the Chemical Weapons Convention, for the verification of these compounds in aqueous samples collected during challenge inspections of suspect production facilities or, in support of allegations of chemical warfare agent use claims.

REFERENCES

1. Ch.E. Kientz, J. Chromatogr.A, 814 (1998) 1.

2. Z. Witkiewicz, M. Mazurek and J. Szulc, J. Chromatogr., 503 (1990) 293.

3. P. A. D'Agostino, L. R. Provost and K. M. Looye, J. Chromatogr., 465 (1989) 271.

4. P. A. D'Agostino, L. R. Provost and J. Visintini, J. Chromatogr., 402 (1987) 221.

5. E. Bonierbale, L. Debordes and L. Coppet, J. Chromatogr. B, 688 (1997) 225.

6. D.K. Rohrbaugh, J. Chromatogr. A, 809 (1998), 131.

7. W. R. Creasy, J. R. Stuff, B. Williams, K. Morrissey, J. Mays, R. Duevel and H. D. Durst, J. Chromatogr. A, 774 (1997) 253.

8. S. Sass and T.L. Fisher, Org. Mass Spectrom., 14 (1979) 257.

9. P. A. D'Agostino and L. R. Provost, Biomed. Environ. Mass Spectrom., 13 (1986) 231.

10. V. T. Borrett, R. J. Mathews and E. R. Mattsson, Aust. J. Chem., 47 (1994) 2065.

11. J.G. Purdon, J.G. Pagotto and R.K. Miller, J. Chromatogr., 475 (1989) 261.

12. P. A. D'Agostino and L. R. Provost, J. Chromatogr., 589 (1992) 287.

13. G. A. Sega, B. A. Tomkins and W. H. Griest, J. Chromatogr. A, 790 (1997) 143.

14. H-A Lakso and W. F. Ng, Anal. Chem., 69 (1997) 1866.

15. C.E. Kientz, A. Verweij, G.J. de Jong and U.A.Th Brinkman, J. Microcol. Sep., 4 (1992) 477.

16. E.R.J. Wils and A.G. Hulst, J. Chromatogr., 454 (1988) 261.

17. E.R.J. Wils and A.G. Hulst, J. Chromatogr., 523 (1990) 151.

18. E.R.J. Wils and A.G. Hulst, Fresenius J. Anal. Chem., 342 (1992) 749.

19. J.A. Tornes, Rapid Commun. Mass Spectrom., 10 (1996) 878.

20. R. M. Black and R.W. Read, J. Chromatogr. A, 759 (1997) 79.

21. R. M. Black and R.W. Read, J. Chromatogr. A, 794 (1998) 233.

22. R. Kostiainen, A.P. Bruins and V.M.A. Hakkinen, J. Chromatogr., 634 (1993) 113.

23. P. A. D'Agostino, L. R. Provost and J. R. Hancock, Mass Spectrometric Identification of Chemical Warfare Agent Precursor and Degradation Products Scheduled by the Chemical Weapons Convention, Proceedings of the 42 nd Annual Conference on Mass Spectrometry and Allied Topics, May 29 - June 3, 1994, Chicago, Illinois, 275.

24. V. T. Borrett, R. Colton and J. C. Traeger, Eur. Mass Spectrom., 1 (1995) 131.

25. V. T. Borrett, R. J. Mathews, R. Colton and J. C. Traeger, Rapid Commun. Mass Spectrom., 10 (1996) 114.

26. P. A. D'Agostino, L. R. Provost and J. R. Hancock, Packed Capillary LC-ESI-MS and MS/MS of CWC Organophosphorus Warfare Agents, Related Compounds and their Degradation Products, Presentation at the 11th Annual Tandem Mass Spectrometry Workshop, Dec. 2-5, 1998, Lake Louise, Alberta, Canada.

27. P. A. D'Agostino, L. R. Provost and J. R. Hancock, J. Chromatogr. A, 808 (1998) 177.

28. Communication at 11th Annual Tandem Mass Spectrometry Workshop with Dr. James Yergey (Merck Frost, Canada).



Figure 1: Packed capillary LC-ESI-MS total-ion-current (600 to 60 u) chromatogram obtained for an aqueous sample containing GB (1), GA (2),GF (3) and GD (4).



Figure 2: Typical ESI-MS mass spectra (sampling cone voltage: 25 volts) acquired for a) GB, b) GA, c) GF and d) GD during LC-ESI-MS analysis.

UNCLASSIFIED

SECURITY CLASSIFICATION OF FORM
(highest classification of Title, Abstract, Keywords)

DOCUMENT CONTROL DATA

(Security classification of title, body of abstract and indexing annotation must be entered when the overall document is classified)

1. ORIGINATOR (the name and address of the organization preparing the document. Organizations for whom the document was prepared, e.g. Establishment sponsoring a contractor's report, or tasking agency, are entered in section 8.)

Defence Research Establishment Suffield
Box 4000, Stn Main
Medicine Hat, AB T1A 8K6 CANADA

2. SECURITY CLASSIFICATION (overall security classification of the document including special warning terms if applicable)

Unclassified

3. TITLE (the complete document title as indicated on the title page. Its classification should be indicated by the appropriate abbreviation (S,C,R or U) in parentheses after the title.)

Analysis of GA, GB, GD and GF in Aqueous Samples by Packed Capillary LC-ESI-MS

4. AUTHORS (Last name, first name, middle initial. If military, show rank, e.g. Doe, Maj. John E.)

D'Agostino, P.A., Hancock, J.R. and Provost, L.R.

5. DATE OF PUBLICATION (month and year of publication of document)

February 1999

6a. NO. OF PAGES (total containing information. Include Annexes, Appendices, etc.)

17

6b. NO. OF REFS (total cited in document)

28

6. DESCRIPTIVE NOTES (the category of the document, e.g. technical report, technical note or memorandum. If appropriate, enter the type of report, e.g. interim, progress, summary, annual or final. Give the inclusive dates when a specific reporting period is covered.)

Suffield Memorandum

8. SPONSORING ACTIVITY (the name of the department project office or laboratory sponsoring the research and development. Include the address.)

9a. PROJECT OR GRANT NO. (if appropriate, the applicable research and development project or grant number under which the document was written. Please specify whether project or grant)

9b. CONTRACT NO. (if appropriate, the applicable number under which the document was written)

10a. ORIGINATOR'S DOCUMENT NUMBER (the official document number by which the document is identified by the originating activity. This number must be unique to this document.)

DRES TM 1999-047

10b. OTHER DOCUMENT NOS. (Any other numbers which may be assigned this document either by the originator or by the sponsor)

11. DOCUMENT AVAILABILITY (any limitations on further dissemination of the document, other than those imposed by security classification)

(X) Unlimited distribution
( ) Distribution limited to defence departments and defence contractors; further distribution only as approved
( ) Distribution limited to defence departments and Canadian defence contractors; further distribution only as approved
( ) Distribution limited to government departments and agencies; further distribution only as approved
( ) Distribution limited to defence departments; further distribution only as approved
( ) Other (please specify):

12. DOCUMENT ANNOUNCEMENT (any limitation to the bibliographic announcement of this document. This will normally correspond to the Document Availability (11). However, where further distribution (beyond the audience specified in 11) is possible, a wider announcement audience may be selected.)

UNCLASSIFIED

SECURITY CLASSIFICATION OF FORM

3. ABSTRACT (a brief and factual summary of the document. It may also appear elsewhere in the body of the document itself. It is highly desirable that the abstract of classified documents be unclassified. Each paragraph of the abstract shall begin with an indication of the security classification of the information in the paragraph (unless the document itself is unclassified) represented as (S), (C), (R), or (U). It is not necessary to include here abstracts in both offical languages unless the text is bilingual).

Packed capillary column liquid chromatography (LC) - electrospray mass spectrometry (ESI-MS) was used for the first time to detect and identify four common organophosphorus chemical warfare agents in aqueous samples. Aqueous samples containing the organophosphorus chemical warfare agents in the 0.01 to 0.1 mg/mL range were analysed directly by packed capillary LC-ESI-MS with the chemical warfare agents and several minor related impurities being well resolved under acetonitrile/water gradient elution conditions. The ESI-MS data for isopropyl methylphosphonofluoridate (GB or sarin), ethyl dimethylphosphoramidocyanidate (GA or tabun), cyclohexyl methylphosphonofluoridate (GF) and pinacolyl methylphosphonofluoridate (GD or soman) were acquired with a sampling cone voltage setting that promoted collisionally activated dissociation, and resulted in the acquisition of informative mass spectra containing both molecular and product ion information. The developed method appears to be an attractive alternative to GC-MS for the analysis of aqueous samples containing organophosphorus chemical warfare agents and their hydrolysis products, since they may be analysed directly without the need for additional sample handling.

4. KEYWORDS, DESCRIPTORS or IDENTIFIERS (technically meaningful terms or short phrases that characterize a document and could be helpful in cataloguing the document. They should be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location may also be included. If possible keywords should be selected from a published thesaurus. e.g. Thesaurus of Engineering and Scientific Terms (TEST) and that thesaurus-identified. If it is not possible to select indexing terms which are Unclassified, the classification of each should be indicated as with the title.)

Mass Spectrometry

Liquid Chromatography

Electrospray

G-Agents

Nerve Agents

Sarin

Soman

Tabun